

Original article

doi: 10.17223/25421379/24/2

GEOCHEMICAL SIMILARITY OF CAMBRIAN ALKALINE AND SUBALKALINE MAGMATISM (KUZNETSK ALATAU OROGEN, SIBERIA): A NEW DATA SYNTHESIS

Vassily V. Vrublevskii¹



¹ National Research Tomsk State University, Tomsk, Russia, vasvr@yandex.ru

Abstract. This is a synthesis of recent geochemical evidence from Cambrian granitic, gabbro-monzonitic, and few small gabbro-foioidolitic intrusions in the Early Paleozoic Kuznetsk Alatau orogen (western Central Asian Orogenic Belt, Siberia). Their origin is attributed to late- and post-orogenic regional magmatism which produced different rock types sharing basic similarity in isotope systematics and trace-element compositions. The $\epsilon_{\text{Nd}}(t)$ values of ~ 2 to 8.7 in main igneous lithologies record the extent of interaction between depleted and enriched mantle. Concurrent increase of initial $^{87}\text{Sr}/^{86}\text{Sr}$ ($0.7039\text{--}0.7058$) ratios and $\delta^{18}\text{O}$ values ($6.5\text{--}13\text{‰}$) in rocks and minerals, as well as ^{207}Pb enrichment ($^{207}\text{Pb}/^{204}\text{Pb}(t) = 15.5\text{--}15.7$), indicate crustal contamination of mantle melts. Trace-element patterns in the rocks correspond to a mixed IAB + OIB source. The alkaline and subalkaline rocks of the Kuznetsk Alatau area share trace element and isotopic similarity and apparently emplaced almost synchronously late during a regional-scale tectonic event. The synthesized data show that Cambrian magmatism in the area involved mixed mantle and continental crust melt components and acted in a complex tectonic setting of the former active margin of the Paleo-Asian Ocean impacted by a mantle plume.

Keywords: Cambrian magmatism, trace-element and isotope geochemistry, mantle plume, continental margin, Paleo-Asian Ocean, Kuznetsk Alatau orogen

Source of financing: this study was supported by the Russian Ministry of Higher Education and Science (Project 0721-2020-0041).

Acknowledgment: I wish to thank my colleagues from the National Research Tomsk State University (Russia) and many other geological institutions of the Russian Academy of Sciences for cooperation in the field and analytical work.

For citation: Vrublevskii V.V. (2022) Geochemical similarity of Cambrian alkaline and subalkaline magmatism (Kuznetsk Alatau Orogen, Siberia): a new data synthesis. *Geosfernye issledovaniya – Geosphere Research*. 3. pp. 27–39. (In Russian). doi: 10.17223/25421379/24/2

Научная статья

УДК 552.11:552.3:550.4(571.17)

doi: 10.17223/25421379/24/2

ГЕОХИМИЧЕСКОЕ СХОДСТВО КЕМБРИЙСКОГО ЩЕЛОЧНОГО И СУБЩЕЛОЧНОГО МАГМАТИЗМА (КУЗНЕЦКИЙ АЛАТАУ, СИБИРЬ): СИНТЕЗ НОВЫХ ДАННЫХ

Василий Васильевич Врублевский¹

¹ Национальный исследовательский Томский государственный университет, Томск, Россия, vasvr@yandex.ru

Аннотация. Обобщены последние геохимические данные по кембрийским гранитоидным, габбро-монзонитовым и габбро-фойдолитовым интрузивам каледонид Кузнецкого Алатау в западной части Центрально-Азиатского орогенического пояса (Сибирь). Их происхождение связано с развитием поздне- и посторогенного регионального магматизма, в результате которого образовались различные изверженные породы, сходные по микроэлементному и изотопному составу. Значения $\epsilon_{\text{Nd}}(t) \sim 2\text{--}8,7$ в главных интрузивных разновидностях отражают степень взаимодействия между деплетированной и обогащенной мантией. Одновременное увеличение первичных отношений $^{87}\text{Sr}/^{86}\text{Sr}$ ($0,7039\text{--}0,7058$) и значений $\delta^{18}\text{O}$ ($6,5\text{--}13\text{‰}$) в породах и минералах, а также обогащение ^{207}Pb ($^{207}\text{Pb}/^{204}\text{Pb}(t) = 15,5\text{--}15,7$) указывают на коровую контаминацию мантийных расплавов. Соотношения редких рассеянных элементов в плутонах соответствуют смешанному IAB + OIB источнику. Щелочные и субщелочные породы Кузнецкого Алатау обладают сходством по микроэлементам и изотопам и, по-видимому, сформировались почти синхронно во время позднего регионального тектогенеза. Синтез данных показывает, что кембрийский магматизм происходил при смешении в расплаве компонентов мантии и континентальной коры в сложной геодинамической обстановке воздействия сублитосферного плюма на ранее сформированную активную окраину Палеоазиатского океана.

Ключевые слова: Кембрийский магматизм, геохимия редких рассеянных элементов и изотопов, мантийный плюм, континентальная окраина, Палеоазиатский океан, Кузнецкий Алатау

Источник финансирования: исследование проведено при поддержке Министерства науки и высшего образования РФ (проект 0721-2020-0041).

Благодарности: автор признателен коллегам из Национального исследовательского Томского государственного университета (Россия) и геологических институтов Российской академии наук за сотрудничество в полевых и аналитических работах.

Для цитирования: Vrublevskii V.V. Geochemical similarity of Cambrian alkaline and subalkaline magmatism (Kuznetsk Alatau Orogen, Siberia): a new data synthesis // Геосферные исследования. 2022. № 3. С. 27–39. doi: 10.17223/25421379/24/2

Introduction

Multiple events of alkaline and subalkaline intraplate magmatism in the geological history of the Central Asian Orogenic Belt (CAOB) have been attributed to the action of mantle plumes [Ernst, 2014; Yarmolyuk, Kuzmin, Ernst, 2014]. Most of Paleozoic intrusions are located in the western flank of the orogenic belt, in the regions of Kuznetsk Alatau, Minusa, Altai-Sayan, Lake Baikal, southeastern Tuva, western Transbaikalia, and northern Mongolia [Yashina, 1982; Nikiforov, Yarmolyuk, 2007; Sklyarov et al., 2009; Yarmolyuk, 2010; Doroshkevich et al., 2012, 2018; Vrublevskii et al., 2012, 2014, 2016a, 2016b, 2018a, 2019, 2020 a, 2020b; Vrublevskii, 2015; Izbrodin et al., 2017; Vrublevskii, Kruk, 2018; Salnikova et al., 2018; Vorontsov et al., 2021; Vrublevskii, Gertner, 2021].

The Early Paleozoic (Caledonian) Kuznetsk Alatau orogen is an island arc terrane within CAOB that formed along the Paleo-Asian active margin. The collisional and accretionary processes were associated with closure of the Paleo-Asian Ocean which, together with the Protopacific ocean, opened as a result of the plume-induced break-up of the Neoproterozoic Rodinia supercontinent about 900–800 Ma ago [Şengör, Natal'in, Burtman, 1993; Khain, 2003; Dobretsov, Buslov, Vernikovskiy, 2003; Windley et al., 2007; Li et al., 2008; Kheraskova et al., 2010; Wilhelm, Windley, 2015; Wan et al., 2018].

The regional-scale event of late- and post-orogenic magmatism, which spanned an interval from ~510 to 490 Ma, produced granitic, gabbro-monzonitic, and mafic alkaline rocks with the respective U-Pb, Sm-Nd, and Rb-Sr ages. It is important to identify the mantle sources of parent melts that produced the diverse subalkaline and alkaline plutonic rocks of the Kuznetsk Alatau terrane [Vrublevskii et al., 2014, 2015, 2016a, 2016b, 2018a].

The aim of this paper is to bring together previously published [Vrublevskii et al., 2016a, 2018a; Vrublevskii, Gertner, 2021] but dispersed geochemical results for specific intrusions within the Kuznetsk Alatau terrane, which are nearly coeval but compositionally different. The synthesized trace-element and isotopic data from

the Cambrian igneous rocks of the area reveal several components in parent magmas that represent an active margin setting and an effect of a mantle plume. The geochemical and geochronological proximity of different igneous lithologies suggests similarity of their magma sources and almost synchronous emplacement in the conditions of a possible plume impact on the lithospheric complexes of a former island arc. The model of regional-scale Cambrian magmatism is new for the Kuznetsk Alatau area. The chronology and evolution trends of the Cambrian magmatism have implications for exchange and/or recycling of material at convergent plate boundaries.

Geological background

The Kuznetsk Alatau accretionary-collisional terrane borders the Minusa and Kuznetsk rift basins in the west and east, respectively (Fig. 1), and has a complex tectonic framework. It comprises abundant Early Paleozoic oceanic and island arc complexes with fragments of the Precambrian basement and small superimposed Devonian graben-like basins [Kungurtsev et al., 2001], as well as deformed Neoproterozoic-Cambrian volcanics and clastic-carbonate sediments, Middle Paleozoic subcontinental volcano-sedimentary deposits, Late Precambrian ophiolites, and Cambrian plutons. Less abundant Devonian and Permian intrusions appear most often as small mafic alkaline stocks and dikes [Vrublevskii, Gertner, 2021]. The general tectonic style of the Kuznetsk Alatau terrane, like the whole Central Asian Orogenic Belt, records the history of its collisions with other terranes during the Paleo-Asian Ocean closure [Dobretsov, Buslov, Vernikovskiy, 2003; Yarmolyuk et al., 2003].

The Cambrian plutons of different compositions discussed in this paper are located in the northern (Mariinsk segment) and eastern (Batenev segment) slopes of the Kuznetsk Alatau Range, where they intrude metamorphosed volcanic and carbonate sediments (Fig. 1 b, 2).

Mafic alkaline magmatism occurs as the Upper Petropavlovka and University small (~0.8–3 km²) gabbroid intrusions cut by foyaitite and few calcite carbonate dikes and veins in the Mariinsk segment.




Fig. 1. (a) Location map of the Kuznetsk Alatau area in the Central Asian Orogenic Belt, simplified after [Şengör, Natal'in, Burtman et al., 1993; Jahn, Wu, Chen, 2000]; (b) tectonic framework of the Kuznetsk Alatau terrane and clusters of Cambrian magmatism, after [Vrublevskii et al., 2016a; Vrublevskii et al., 2018a; Vrublevskii, Gertner, 2021]

Рис. 1. (a) Схема размещения Кузнецкого Алатау в Центрально-Азиатском орогеническом поясе, по [Şengör, Natal'in, Burtman et al., 1993; Jahn, Wu, Chen, 2000]; (b) тектоническая схема террейна Кузнецкого Алатау и проявлений кембрийского магматизма, по [Vrublevskii et al., 2016a; Vrublevskii et al., 2018a; Vrublevskii, Gertner, 2021]

The foidolite and carbonatite rocks have Sm-Nd isochron ages of ~510–500 Ma [Vrublevskii, 2015; Mustafaev et al., 2020]. Gabbro-monzonite plutons of similar ages (~490–505 Ma, U-Pb) cluster among Early Paleozoic granitoids of the Batenev segment. The structural and compositional frameworks of large (~40 to 200 km²) plutons correspond to a two-phase history [Dovgal', Shirokikh, 1980; Vrublevskii et al., 2018a]. The Cambrian (~490–510 Ma, U-Pb) plutons reaching sizes of 60–70 to 500 km² consist mainly of granodiorite-tonalite and later granites [Vrublevskii et al., 2016a]. Coeval granitoids in the Mariinsk segment form up to 300–500 km² batholiths composed of diorite-tonalite, plagiogranite, and granite, as well as quartz syenite and granosyenite [Rudnev, 2013].

Compositions of main igneous lithologies

Various aspects of chemistry and isotope systematics of the Kuznetsk Alatau igneous rocks were detailed in numerous earlier publications [Rudnev, 2013; Vrublevskii, 2015; Vrublevskii et al., 2016a, 2018a; Mustafaev et al., 2020; Vrublevskii, Gertner, 2021].

Main features of different rock types are summarized as selected analyses in Tables 1 and 2 and illustrated by trace-element and Nd and Sr isotope patterns in Figures 3 and 4.

Major- and trace-element chemistry.

The Kuznetsk Alatau *gabbro-foidolite* intrusions typically have low silica (~44–53 wt. % SiO₂), high calcium (to ~10–15 wt. % CaO), and elevated aluminum (to 18–22 wt. % Al₂O₃) and alkali (to 9–12 wt. % (Na₂O + K₂O), Na₂O/K₂O ≈ 2–7) contents. Compared to gabbro, the later foyaite and foidolite have lower MgO# (0.5–0.1) and compatible elements (529–4 ppm Cr, 112–3 ppm Ni, 200–8 ppm V, 48–2 ppm Co), but higher LILE and HFSE reaching ~10 to 90 ppm Rb, ~2000 ppm Ba, 540 to 1180 ppm Sr, 0.4 to 7 ppm Th, 0.3 to 5.4 ppm U, 46 to 240 ppm REE, ~100 to 280 ppm Zr, and ~10 to 50 ppm Nb). The patterns of REE (La/Yb_N ≈ 6–11; Eu/Eu* ≈ 1) and other trace elements in these intrusions look like those of average ocean island basalts (OIB) but bear additional island-arc basalt (IAB) signatures. The IAB contribution appears in multi-element spectra with distinct Nb-Ta and Zr-Hf depletion.

The *gabbro-monzonitic* intrusions of the area contain moderate amounts of alkalis (~3–10 wt. % $\text{Na}_2\text{O} + \text{K}_2\text{O}$) and show large ranges of silica (~43 to 65 wt. % SiO_2) and alumina (~15–20 wt. % Al_2O_3), while Fe# is about 0.5. Some of the rocks have relatively high K_2O contents of ~5–6 wt. % ($\text{K}_2\text{O}/\text{Na}_2\text{O}$ from 0.2–0.7 to 0.5–1.5), along with quite low TiO_2 (~2.2 to 0.5 wt. %). The concentrations of compatible elements decrease markedly in the series from gabbro to monzodiorite and monzonite: 436 to 12 ppm Cr, 182 to 9 ppm Ni, 574 to 32 ppm V, and 56 to 5 ppm Co,

whereas LILE and HFSE become times higher: ~3 to 134 ppm Rb, ~100 to 2900 ppm Ba, ~0.2 to 31 ppm Th, ~0.4 to 37 ppm Nb, and ~60 to 336 ppm REE ($\text{Eu}/\text{Eu}^* \approx 0.7\text{--}1.3$); Sr reaches ~1400 to 2500 ppm. The REE ($\text{La}/\text{Yb}_N \approx 7\text{--}30$) and other trace-element patterns in gabbro are similar to the average IAB composition, with typical minimums in Nb-Ta and Zr-Hf. The LILE and HFSE concentrations in more strongly fractionated ($\text{La}/\text{Yb}_N \approx 14\text{--}34$) monzonites approach or even exceed the OIB average but the general shapes of the spectra remain as in IAB.




Fig. 2. (a) Location map of gabbro-foioidlite plutons (not to scale) and (b) granitoids in the Mariinsk segment, after [Rudnev, 2013; Vrublevskii et al., 2014; Vrublevskii, Gertner, 2021]

Gabbro-foioidlite intrusions: UP = Upper Petropavlovka, Un = University; (b) Geological framework of gabbro-monzonitic and granitic intrusions in the Batenev segment, after [Dovgal', Shirokikh, 1980; Vrublevskii et al., 2016a, 2018a]. Gabbro-monzonite intrusions: Kg = Kogtakh, Bl = Balakhcha, Ksh = Kashpar, Ks = Kiskach, Cht = Chas-Taiga, Pt = Pistag, Kt = Karatag

Рис. 2. (а) Локализация габбро-фойдолитовых плутонов (вне масштаба) и (b) гранитоидов в Мариинском сегменте, по [Vrublevskii et al., 2014; Vrublevskii, Gertner, 2021; Rudnev, 2013]

Габбро-фойдолитовые интрузивы: UP = Верхнепетропавловский, Un = Университетский; (b) Геологическая карта габбро-монзонитовых и гранитных интрузивов в Батеневском сегменте, по [Dovgal', Shirokikh, 1980; Vrublevskii et al., 2016a, 2018a]. Габбро-монзонитовые интрузивы: Kg = Когтахский, Bl = Балахчинский, Ksh = Кашпарский, Ks = Кискачинский, Cht = Частайгинский, Pt = Пистагский, Kt = Каратагский

Table 1

Selected analyses of the Kuznetsk Alatau Cambrian granitic, gabbro-monzonitic, and gabbro-foiolitic rocks

Таблица 1

Выборка анализов кембрийских гранитоидов, габбро-монзонитов и габбро-фойдолитов Кузнецкого Алатау

Intrusion	Gabbro-foiolite			Gabbro-monzonite			Granitoids					
Rock type	SG	AG	I	SG	MD	M	Tn	Tn	GD	S	GS	Gr
SiO ₂ , wt, %	44,98	46,46	44,38	44,77	53,47	58,90	64,04	65,72	67,85	63,14	68,60	70,76
TiO ₂	0,95	1,27	0,91	1,57	0,89	1,01	0,43	0,41	0,32	0,60	0,32	0,17
Al ₂ O ₃	15,11	14,71	18,57	14,31	18,89	17,59	16,28	15,09	15,26	17,26	16,65	15,54
Fe ₂ O ₃	11,20	11,34	10,68	14,83	7,83	6,51	5,00	5,82	3,84	3,69	2,88	2,03
MnO	—	—	0,22	0,14	0,12	0,10	0,12	0,10	0,11	0,08	0,06	0,08
MgO	8,93	6,92	2,34	8,12	4,42	3,41	4,12	2,33	1,33	0,51	0,81	0,76
CaO	14,63	10,53	11,42	12,52	7,44	4,67	3,01	4,24	2,46	3,15	2,55	1,35
Na ₂ O	2,96	4,23	7,22	2,25	4,55	4,06	4,53	1,65	5,07	7,42	4,19	5,55
K ₂ O	0,95	2,43	2,44	0,96	1,96	2,58	1,94	1,64	2,25	3,38	3,20	3,35
P ₂ O ₅	0,09	0,50	0,69	0,21	0,53	0,33	0,07	0,09	0,05	0,22	0,05	0,06
LOI	1,13	1,26	1,18	0,41	0,13	0,30	0,43	1,82	0,71	0,24	0,12	0,17
Total	100,93	99,65	100,05	100,09	100,23	99,46	99,98	98,91	99,25	99,70	99,43	99,82
Cr, ppm	224	56	28	57	49	35	96	—	51	38	—	40
Ni	56	7	22	86	16	15	32	—	16	7	—	8
V	155	12	81	274	159	125	—	—	—	—	—	—
Co	49	12	21	56	26	17	15	—	10	4	—	5
Sc	24	0,9	5,8	53	19	12	11	—	6	1,6	—	2,4
Pb	—	—	5,4	—	13	11	7	—	9	11	—	14
Cs	0,8	1	0,5	0,6	0,7	1,6	0,9	—	0,4	0,2	—	0,8
Rb	24	42	34	13	69	83	51	44	62	94	64	85
Ba	303	726	1802	231	1105	1076	658	531	1024	1110	1030	785
Sr	538	893	869	720	1407	1336	648	401	809	979	442	745
Nb	9	42	12	4,6	19	21	11	2,4	19	16	14	22
Ta	0,6	2,5	0,7	0,24	1,1	1,5	0,8	0,4	1,4	1	1,3	1,7
Zr	124	279	144	71	157	368	23	23	47	76	209	42
Hf	2,7	4	2,6	1,9	2,6	4,7	0,8	1,2	1,6	2,2	6,5	1,4
Y	22	45	27	15	31	18	10	19	15	11	10	11
Th	2,7	7	3	1,7	6,7	8,9	2,7	2,2	7,5	4,2	4,9	5
U	1,9	5,5	2	0,53	1,5	1,8	1,2	0,8	1,3	0,8	2,6	1,8
La	21	49	32	13	52	49	22	11	47	31	31	35
Ce	45	101	63	29	115	125	43	23	84	60	52	63
Pr	5,2	9,1	7,3	4,4	13	14	5	2,2	8,3	6	6,2	6,4
Nd	21	44	28	20	45	55	19	10	30	20	20	24
Sm	4,6	8,5	5,3	3,9	7,4	9,8	3,3	2,2	4,7	3,1	2,9	3,7
Eu	1,3	2,6	1,8	1,2	2,4	2,3	1	0,72	1,2	0,75	1	0,87
Gd	4,4	8,1	4,8	3,6	6,6	9,1	3,2	2,3	4,5	2,4	2,2	3,4
Tb	0,7	1,3	0,7	0,51	0,8	1,2	0,45	0,4	0,63	0,37	0,36	0,5
Dy	4,3	8,2	4,2	2,9	3,8	6,4	2,1	2,6	3	1,7	2	2,3
Ho	0,9	1,8	0,9	0,57	0,77	1,3	0,4	0,6	0,57	0,35	0,4	0,43
Er	2,4	5,1	2,5	1,5	2,1	3,4	1,1	1,9	1,6	0,98	1,1	1,2
Tm	0,4	0,8	0,4	0,2	0,3	0,53	0,15	0,35	0,22	0,15	0,21	0,17
Yb	2,2	4,8	2,6	1,3	1,8	2,3	1	2,1	1,6	1	1,4	1,1
Lu	0,3	0,7	0,4	0,18	0,27	0,35	0,15	0,29	0,24	0,16	0,18	0,17
ΣREE	114	245	155	82	251	280	102	59	188	128	121	142
LREE/HREE	6,3	7	8,3	6,8	14,5	10,2	10,7	5	13,7	16,1	14,3	14,7

Note. SG = subalkaline gabbro, AG = alkaline gabbro, I = feldspar ijolite, MD = monzodiorite, M = monzonite, Tn = tonalite, GD = granodiorite, S = quartz syenite, GS = granosyenite, Gr = subalkaline granite. (—) is not detected. Major and trace elements and REE in rocks were analyzed by XRF and ICP–MS, respectively, under standard operating conditions. Data after [Rudnev, 2013; Vrublevskii, 2015; Vrublevskii et al., 2016a, 2018a; Mustafaev et al., 2020].

Примечания. SG = субщелочное габбро, AG = щелочное габбро, I = полевошпатовый ийолит, MD = монцодиорит, M = монзонит, Tn = тоналит, GD = гранодиорит, S = кварцевый сиенит, GS = граносиенит, Gr = субщелочной гранит. (—) не обнаружено. Петрогенные, редкие и редкоземельные элементы в породах анализировались методами XRF и ICP–MS соответственно, в стандартных условиях. Данные по [Rudnev, 2013; Vrublevskii, 2015; Vrublevskii et al., 2016a, 2018a; Mustafaev et al., 2020].




Fig. 3. Chemical classification and compositions of the Kuznetsk Alatau Cambrian igneous rocks

a – R1–R2 diagram [De la Roche et al., 1980]; **b–d** – Primitive mantle-normalized [Sun, McDonough, 1989] multi-element spectra. All diagrams include selected analyses (Table 1) and published data from [Rudnev, 2013; Vrublevskii, 2015; Vrublevskii et al., 2016a, 2018a; Mustafaev et al., 2020; Vrublevskii, Gertner, 2021]. Average ocean island basalt (OIB) and island-arc basalt (IAB) compositions are after [Sun, McDonough, 1989; Kelemen, Hanghøj, Greene, 2003], respectively

Рис. 3. Химическая классификация и состав кембрийских изверженных пород Кузнецкого Алатау

a – диаграмма R1–R2 [De la Roche et al., 1980]; **b–d** – мультиэлементные спектры (нормализация по составу примитивной мантии [Sun, McDonough, 1989]). На диаграммах нанесены табличные (табл. 1) и литературные данные по [Rudnev, 2013; Vrublevskii, 2015; Vrublevskii et al., 2016a, 2018a; Mustafaev et al., 2020; Vrublevskii, Gertner, 2021]. Средние составы базальтов океанских островов (ocean island basalt = OIB) и островных дуг (island-arc basalt = IAB) по [Sun, McDonough, 1989; Kelemen, Hanghøj, Greene, 2003] соответственно

Most of the Kuznetsk Alatau granitoids are subalkaline, alkaline or less often peralkaline varieties (~58–76 wt. % SiO₂; ~3–13 wt. % (Na₂O + K₂O); K₂O/Na₂O ~0.2–1.7) with a relatively low Al saturation index (ASI ≤1.1; suprasubduction granitoids) and Fe# ≤0.8. The REE concentrations are quite low (~50–230 ppm; La/Yb_N ~ from 3–8 to 15–37) while the HFSE distribution corresponds to the average IAB composition with Nb-Ta and Zr-Hf minima. Some granitoids show above-OIB enrichment in several trace elements that reach ~140 ppm Rb, ~1110 ppm Ba, ~1100 ppm Sr, ~8 ppm U, and ~40 ppm Th.

Isotope systematics (Nd-Sr-Pb-O).

The gabbro-foidolite intrusions of the area show notable variations of initial Nd and Sr isotope ratios, ¹⁴³Nd/¹⁴⁴Nd(t) = 0.512245–0.512438 (ε_{Nd}(t) ~5–8.7) and ⁸⁷Sr/⁸⁶Sr(t) = 0.7041–0.7058, as well as the ~7–13 ‰

^δ¹⁸O_{V-SMOW} values and the ²⁰⁷Pb enrichment (²⁰⁷Pb/²⁰⁴Pb(t) 15.53–15.71, ²⁰⁶Pb/²⁰⁴Pb(t) 17.92–20.65) in rocks and minerals [Vrublevskii, Gertner, 2021].

The gabbro-monzonitic plutons have less radiogenic Nd and Sr isotope compositions (¹⁴³Nd/¹⁴⁴Nd(t) = 0.512174–0.512273; ε_{Nd}(t) ~3.5–5.4; ⁸⁷Sr/⁸⁶Sr(t) = 0.7039–0.7052). Their Nd isotope heterogeneity is more prominent than in the gabbro-foidolite rocks. On the other hand, the ⁸⁷Sr/⁸⁶Sr and ^δ¹⁸O_{V-SMOW} (6.5–8.8 ‰) values are concordantly high [Vrublevskii et al., 2018a]. The granitoids of the area share isotopic similarity with the gabbro-monzonitic rocks: ¹⁴³Nd/¹⁴⁴Nd(t) = 0.512098–0.512346; ε_{Nd}(t) ~2–4.8; ⁸⁷Sr/⁸⁶Sr(t) ~0.7047–0.7052 [Rudnev, 2013; Vrublevskii et al., 2016a]. The isotopic heterogeneity of the igneous rock associations may record participation of different magma sources in the parent melts.

Table 2

Sr-Nd isotopic composition of the Kuznetsk Alatau Cambrian granitic, gabbro-monzonitic, and gabbro-foidolitic rocks


Таблица 2

Sr-Nd изотопный состав кембрийских гранитоидов, габбро-монзонитов и габбро-фойдолитов Кузнецкого Алатау

Rock series	Rock-type	Sm, ppm	Nd, ppm	$^{143}\text{Nd}/^{144}\text{Nd}$	$^{143}\text{Nd}/^{144}\text{Nd}(t)$	$\epsilon_{\text{Nd}}(t)$	Rb, ppm	Sr, ppm	$^{87}\text{Sr}/^{86}\text{Sr}$	$^{87}\text{Sr}/^{86}\text{Sr}(t)$
Gabbro-foidolitic	SG	3,42	15,3	0,512808	0,512365	7,25	19	583	0,70620	0,70553
	“	1,77	7,46	0,512907	0,512438	8,67	14	745	0,70520	0,70483
	“	2,74	9,43	0,512937	0,512362	7,19	8,6	709	0,70452	0,70414
	AG	5,98	31,38	0,512766	0,512389	7,71	44	964	0,70649	0,70557
	NS	9,45	50,77	0,512727	0,512432	6,03	207	539	0,71266	0,70495
	I	5,18	27,5	0,512618	0,512245	4,90	19	1365	0,70566	0,70538
	C	2,97	11,9	0,512740	0,512247	4,94	3	1000	0,70590	0,70584
Gabbro-monzonitic	SG	7,84	36,90	0,512627	0,512206	4,14	7,24	1626	0,70414	0,70405
	“	4,91	26,67	0,512538	0,512174	3,52	51,2	961	0,70576	0,70469
	“	9,15	41,78	0,512661	0,512227	4,55	9,7	1164	0,70539	0,70522
	MD	7,88	46,93	0,512511	0,512179	3,61	72,3	1278	0,70551	0,70437
	“	6,21	38,78	0,512513	0,512196	3,95	45,1	1179	0,70561	0,70484
	“	10,79	63,43	0,512538	0,512201	4,04	52,5	2180	0,70450	0,70402
	M	6,27	36,04	0,512586	0,512241	4,82	67,3	733	0,70704	0,70519
	“	3,13	21,02	0,512568	0,512273	5,45	62,8	1092	0,70587	0,70472
	“	6,08	40,36	0,512494	0,512196	3,95	41,3	3073	0,70413	0,70386
Granitic	GD	4,09	24,5	0,512563	0,512226	4,83	42,4	886	0,70576	0,70478
	S	3,88	24,9	0,512543	0,512240	4,59	44,6	1118	0,70557	0,70478
	Gr	3,65	22,9	0,512429	0,512121	2,26	73,2	804	0,70693	0,70514

Note. NS = nepheline syenite, C = carbonatite. Isotopic data after [Vrublevskii et al., 2016a, 2018a; Mustafaev et al., 2020; Vrublevskii, Gertner, 2021]. The isotope analyses were carried out using the standard techniques and instruments.

Примечания. NS = нефелиновый сиенит, C = карбонатит. Данные по [Vrublevskii et al., 2016a, 2018a; Mustafaev et al., 2020; Vrublevskii, Gertner, 2021]. Изотопные анализы проводились с использованием стандартных методик и оборудования.

Fig. 4. $\epsilon_{\text{Nd}}(t)$ vs. $^{87}\text{Sr}/^{86}\text{Sr}(t)$ plot for the Kuznetsk Alatau Cambrian igneous rocks

Average basalt compositions are after [Yarmolyuk, Kovalenko, 2003] for the Altai-Sayan fold area (NA = North Asian plume) and [Lightfoot et al., 1993] for the Siberian Trap Province (S = Siberian plume). Mantle array, DMM (depleted MORB mantle), PREMA (prevalent mantle), EM 1 (enriched mantle 1) and EM 2 (enriched mantle 2) mantle reservoirs are according to [Zindler, Hart, 1986; Salters, Stracke, 2004]

Рис. 4. Диаграмма $\epsilon_{\text{Nd}}(t)$ – $^{87}\text{Sr}/^{86}\text{Sr}(t)$ для кембрийских изверженных пород Кузнецкого Алатау

NA (North Asian plume) = средний состав базальтов Алтае-Саянской складчатой области, по [Yarmolyuk, Kovalenko, 2003] S (Siberian plume) = средний состав базальтов Сибирской трапповой провинции, по [Lightfoot et al., 1993]. Мантийная последовательность (Mantle array), резервуары DMM (depleted MORB mantle), PREMA (prevalent mantle), EM 1 (enriched mantle 1) and EM 2 (enriched mantle 2) мантии, по [Zindler, Hart, 1986; Salters, Stracke, 2004]

Discussion

Major-element chemistry of Cambrian intrusions and its implications.

The discussed plutonic complexes differ markedly in petrography and major-element chemistry. The low-silica (~43–47 wt. % SiO₂) rocks have mainly sodic alkalinity (~0.9–6.5 wt. % Na₂O + K₂O, Na₂O/K₂O ≥ 1), which is common to rift-related mantle alkaline magmatism. The higher-SiO₂ (~58–76 wt. %) varieties, including granitoids and most of monzonites, have higher total alkalis (~10–13 wt. % Na₂O + K₂O) and variable K₂O/Na₂O ratios from ~0.2 to 1.7. The K₂O contents in some monzonite samples are relatively high (~5–6 wt. % and K₂O/Na₂O up to ~1.5) and, along with low TiO₂ (~0.3–2.2 wt. %), may represent continental-margin material at the source. The Al₂O₃ contents of ~12–18 wt. % and relatively low ASI values (≤1.1) in most of the granitoids correspond to *I*-type granite compositions inconsistent with purely collisional environments. Thus, the observed whole-rock chemistry features provide implicit evidence for a complex tectonic setting of magmatism.

Sources of Cambrian plutonic magmatism in the Kuznetsk Alatau orogen

The Cambrian (~510–490 Ma) gabbro-foiolitic, gabbro-monzonitic, and granitic intrusions in the Kuznetsk Alatau orogen emplaced synchronously with magmatic events in the Western CAO Early Paleozoic large igneous province (LIP) [Izokh et al., 2008; Vrublevskii et al., 2012; Yarmolyuk, Kuzmin, Ernst, 2014]. The Nd isotope composition of rocks reveals a mantle contribution to the parent magma generation. The large $\epsilon_{\text{Nd}}(t)$ range from 8.7 to 2 may result from mixing and homogenization of the depleted (PREMA-like) and enriched (EM-like) lithospheric mantle components.

The gabbro-foiolite rocks are derived from the most depleted mantle, while the share of the EM component is greater in the gabbro-monzonitic and granitic rocks (Fig. 4). Isotopic heterogeneity of this kind in roughly coeval igneous complexes is possible within even a small field of magmatism [Bell, Tilton, 2001]. A similar $\epsilon_{\text{Nd}}(t)$ range of ~1 to 7 is common to Cambrian alkaline plutons elsewhere in the LIP (Russian Altai, SE Tuva, Western Transbaikalia), which have a proven mantle origin [Doroshkevich et al., 2012; Vrublevskii et al., 2012, 2020a]. These Kuznetsk Alatau intrusions, as well as other occurrences of Early Paleozoic magmatism in the Altai-Sayan region, may result from the activity of the North Asian PREMA superplume [Yarmolyuk, Kovalenko, 2003; Kuzmin, Yarmolyuk, Kravchinsky, 2010].

On the other hand, the concurrently increasing $^{87}\text{Sr}/^{86}\text{Sr}(t) = 0.7039\text{--}0.7058$ (Fig. 4) and $\delta^{18}\text{O}_{\text{SMOW}} \sim 6.5\text{--}13\text{‰}$ values in the Kuznetsk Alatau igneous rocks may result from crustal contamination of the parental

melts. The interaction of the melts with continental crust is further supported by enrichment in radiogenic lead, with the initial $^{207}\text{Pb}/^{204}\text{Pb}$ 15.71 and $^{206}\text{Pb}/^{204}\text{Pb}$ 20.65 ratios typical of those in orogenic areas [Vrublevskii, Gertner, 2021]. Note that many Paleozoic-Mesozoic volcano-plutonic complexes in the Western CAO share signatures of additional upper crustal ^{87}Sr and ^{18}O inputs [Pokrovskii et al., 1998; Nikiforov et al., 2002; Vrublevskii et al., 2003, 2005, 2012, 2015, 2016a, 2016b, 2018a, 2018b, 2019, 2020a, 2020b; Doroshkevich et al., 2012, 2018; Baatar et al., 2013; Krupchatnikov, Vrublevskii, Kruk, 2018; Vrublevskii, Gertner, Chugaev, 2018; Vrublevskii, Gertner, 2021].

Post-accretion continental margin magmatism under a plume impact.

Despite the enrichment in some LILEs and HFSEs till the OIB level, the trace-element patterns of gabbro-foiolitic, gabbro-monzonitic, and granitic rocks generally show IAB affinity (Fig. 3 b–d). Therefore, magma generation involved mixed OIB-like (within-plate plume) and IAB-like (subduction) components. This mixing is consistent with the idea that the Kuznetsk Alatau accretionary orogen evolved on the former active margin of the Paleo-Asian Ocean [Kungurtsev et al., 2001; Dobretsov, Buslov, Vernikovskiy, 2003], which was exposed to the impact of a plume.

Different contributions of the IAB and OIB components to the melts show up in LILE and HFSE variations observed in the analyzed samples, with Th, Ta, Nb, Yb, and Y ranges corresponding to OIB/E-MORB-type sources of enriched mantle (Fig. 5 a–b, f). At the same time, island arc material includes continental crust components, which may be partly responsible for the degree of Nb depletion and its correlation with Zr, as well as for the U, Ba, Rb, and Sr enrichment of the Kuznetsk Alatau igneous rocks (Fig. 3, 5 c–e). Although the $\epsilon_{\text{Nd}}(t)$ values are mostly in the ~3–6 range typical of rocks derived from moderately depleted mantle, the Ce/Pb, Ce/Nb, and Th/Nb ratios vary markedly (Fig. 4, 6), possibly, because the mantle melts were more or less strongly contaminated with island arc material.

Thus, the trace-element patterns of the Kuznetsk Alatau gabbro-foiolitic, gabbro-monzonitic, and granitic rocks record the source heterogeneity, as well as a complex tectonic setting of their emplacement. The geochemical similarity of the Cambrian intrusions in the Kuznetsk Alatau orogen may be due to Early Paleozoic activity of the PREMA/OIB-superplume [Yarmolyuk, Kovalenko, 2003], which impinged on the lithosphere beneath the former active margin of the Paleo-Asian Ocean.

The $\epsilon_{\text{Nd}}(t)$ behavior records variations of depleted and enriched material in the parent mantle domain and is concordant with the general isotopic evolution trend of magma sources from the Early Paleozoic (North Asian plume) through earliest Mesozoic (Siberian plume) time (Fig. 4).




Fig. 5. Trace-element contents and their ratios in the Kuznetsk Alatau Cambrian igneous rocks

a – Th/Yb–Ta/Yb diagram [Gorton, Schandl, 2000]: OIA = oceanic island arc, ACM = active continental margin, WPVZ = within-plate volcanic zone, E-MORB = enriched mid-ocean ridge basalt; WPB = within-plate basalt; **b** – ThN–NbN diagram [Saccani, 2015]: AB = alkaline basalt, BABB = back-arc basin basalt; N-MORB-normalized [Sun, McDonough, 1989] Th and Nb; **c** – Ba/Nb–La/Nb diagram [Bi et al., 2015]; **d** – Rb–(Y + Nb) diagram [Pearce, Harris, Tindle, 1984]: syn-COLG = syncollisional granite, VAG = volcanic arc granite; WPG = within plate granite; **e** – (Nb/Zr)_N–Zr diagram [Thièblemont, 1999]: BSE-normalized [Hofmann, 1988] Nb/Zr; **f** – Yb/Ta–Y/Nb diagram [Eby, 1990]. Diagrams in panels d and e include only granitic and monzonitic compositions

Рис. 5. Соотношения микроэлементов в кембрийских изверженных породах Кузнецкого Алатау

a – диаграмма Th/Yb–Ta/Yb [Gorton, Schandl, 2000]: OIA = океанская островная дуга (oceanic island arc), ACM = активная континентальная окраина (active continental margin), WPVZ = внутриплитная вулканическая зона (within-plate volcanic zone), E-MORB = обогащенный базальт срединно-океанического хребта (enriched mid-ocean ridge basalt); WPB = внутриплитный базальт (within-plate basalt); **b** – диаграмма ThN–NbN [Saccani, 2015]: AB = щелочной базальт (alkaline basalt), BABB = базальт задугового бассейна (back-arc basin basalt); содержание Th и Nb нормализовано по N-MORB [Sun, McDonough, 1989]; **c** – диаграмма Ba/Nb–La/Nb [Bi et al., 2015]; **d** – диаграмма Rb–(Y + Nb) [Pearce, Harris, Tindle, 1984]: syn-COLG = синколлизионный гранит (syncollisional granite), VAG = островодужный гранит (volcanic arc granite); WPG = внутриплитный гранит (within plate granite); **e** – диаграмма (Nb/Zr)_N–Zr [Thièblemont, 1999]: Значения Nb/Zr нормализованы по BSE (Bulk Silicate Earth) [Hofmann, 1988]; **f** – диаграмма Yb/Ta–Y/Nb [Eby, 1990]. На панелях «d» и «e» нанесены составы только гранитов и монзонитов




Fig. 6. Ce/Nb–Th/Nb (a) and $\epsilon_{\text{Nd}}(t)$ –Ce/Pb (b) diagrams for the Kuznetsk Alatau Cambrian igneous rocks
Compositions of primitive (PM) and depleted (DMM) mantle, E-MORB, OIB, PREMA, BSE (bulk silicate Earth), and UC (upper crust) are after [Zindler, Hart, 1986; Sun, McDonough, 1989; Rudnick, Gao, 2003; Salters, Stracke, 2004]

Рис. 6. Диаграммы Ce/Nb–Th/Nb (a) и $\epsilon_{\text{Nd}}(t)$ –Ce/Pb (b) для кембрийских изверженных пород Кузнецкого Алатау

Составы примитивной (PM) и деплетированной (DMM) мантии, E-MORB, OIB, PREMA, BSE и UC (upper crust = верхняя кора), по [Zindler, Hart, 1986; Sun, McDonough, 1989; Rudnick, Gao, 2003; Salters, Stracke, 2004]

Meanwhile, some trace-element and isotopic dissimilarity of the rocks may result from their emplacement among compositionally and structurally heterogeneous accretionary-collisional complexes. Signatures of such plume-lithosphere interaction were reported from different areas of the Paleo-Asian active margins [Dobretsov, 2011; Vrublevskii et al., 2012, 2016a, 2018a; Gordienko, Metelkin, 2016; Gordienko, 2019; Vrublevskii, Gertner, 2021].

The subcontinental lithospheric mantle (SCLM) apparently played a subordinate role in the process, as a mantle wedge effect. Any significant involvement of SCLM components was hardly possible, given the island arc origin of the Kuznetsk Alatau terrane and the Nd isotope composition of the analyzed rocks corresponding to moderately depleted mantle.

Conclusion

Cambrian magmatism in the Kuznetsk Alatau orogen produced granitic, gabbro-monzonitic, and gabbro-

foidolitic intrusions within the ~510 to 490 Ma time span and terminated the collisional processes along the Early Paleozoic Paleo-Asian active margin. The observed Sm–Nd isotopic patterns in main igneous lithologies indicate parentage of the magma sources and involvement of PREMA-type and EM-type mantle components in the magma generation. The existence of the prevalent mantle (PREMA) reservoir was possibly maintained by the activity of the North Asian superplume, which induced the Early Paleozoic magmatism of the Western CAOB.

The trace-element compositions of igneous rocks record possible mixing of IAB- and OIB-like components in the source magma, whereas relatively high Sr, Pb and O isotope ratios record crustal contamination of the melts. The revealed geochemical similarity of the Kuznetsk Alatau Cambrian igneous complexes is not fortuitous but may be due to their emplacement in the former active margin exposed to the impact of a mantle plume.

References

- Baatar M., Ochir G., Kynicky J., Iizumi S., Comin-Chiaramonti P. Some notes on the Lugin Gol, Mushgai Khudag and Bayan Khoshuu alkaline complexes, Southern Mongolia // *International Journal of Geosciences*. 2013, 4, pp.1200–1214.
- Bell K., Tilton G.R. Nd, Pb and Sr isotopic compositions of East African carbonatites: Evidence for mantle mixing and plume inhomogeneity // *Journal of Petrology*. 2001, 42, pp. 1927–1945.
- Bi J.H., Ge W.C., Yang H., Zhao G.C., Xu W.L., Wang Z.H. Geochronology, geochemistry and zircon Hf isotopes of the Dongfanghong gabbroic complex at the eastern margin of the Jiamusi Massif, NE China: petrogenesis and tectonic implications // *Lithos*. 2015, 234–235, pp. 27–46.
- De la Roche H., Leterrier J., Grandclaude P., Marchal M. A classification of volcanic and plutonic rocks using R_1 – R_2 diagram and major element analyses – Its relationships with current nomenclature // *Chemical Geology* 1980, 29, pp. 183–210.
- Dobretsov N.L. Early Paleozoic tectonics and geodynamics of Central Asia: role of mantle plumes // *Russian Geology and Geophysics*. 2011, 52, pp. 1539–1552.

- Dobretsov N.L., Buslov M.M., Vernikovskiy V.A. Neoproterozoic to Early Ordovician evolution of the Paleo-Asian ocean: implications to the break-up of Rodini // *Gondwana Research*. 2003, 6, pp. 143–159.
- Doroshkevich A.G., Izbrodin I.A., Rampilov M.O., Ripp G.S., Lastochkin E.I., Khubanov V.B. Permo–Triassic stage of alkaline magmatism in the Vitim plateau (western Transbaikalia) // *Russian Geology and Geophysics*. 2018, 59, pp. 1061–1077. DOI 10.1016/j.rgg.2018.08.001.
- Doroshkevich A.G., Ripp G.S., Izbrodin I.A., Savatenkov V.M. Alkaline magmatism of the Vitim province, West Transbaikalia, Russia: Age, mineralogical, geochemical and isotope (O, C, D, Sr and Nd) data // *Lithos*. 2012, 152, pp.157–172.
- Dovgal' V.N., Shirokikh V.A. *Istoriya razvitiya magmatizma povyshennoy shchelochnosti Kuznetskogo Alatau* [Evolution of High-Alkalinity Magmatism in the Kuznetsk Alatau], Novosibirsk: Nauka, 1980, 215 p. In Russian: Довгаль В.Н., Широких В.А. История развития магматизма повышенной щелочности Кузнецкого Алатау. Новосибирск: Наука. 1980. 215 с.
- Eby G.N. The A-type granitoids: a review of their occurrence and chemical characteristics and speculation on their petrogenesis. *Lithos* 1990, 26, pp. 115–134.
- Ernst R.E. Large Igneous Provinces, Cambridge University Press: Cambridge, 2014. 630 p.
- Gordienko I.V. Relationship between subduction-related and plume magmatism at the active boundaries of lithospheric plates in the interaction zone of the Siberian continent and Paleozoic ocean in the Neoproterozoic and Paleozoic // *Geodynamics and Tectonophysics*. 2019, 10, pp. 405–457.
- Gordienko I.V., Metelkin D.V. The evolution of the subduction zone magmatism on the Neoproterozoic and Early Paleozoic active margins of the Paleo Asian ocean // *Russian Geology and Geophysics*. 2016, 57, pp. 69–81.
- Gorton M.P., Schandl E.S. From continents to island arcs: a geochemical index of tectonic setting for arc-related and within-plate felsic to intermediate volcanic rocks // *The Canadian Mineralogist*. 2000, 38, pp. 1065–1073, DOI 10.2113/gscanmin.38.5.1065.
- Hofmann A.W. Chemical differentiation of the Earth. The relationship between mantle, continental crust and oceanic crust // *Earth and Planetary Science Letters*. 1988, 90, pp. 297–314.
- Izbrodin I.A., Doroshkevich A.G., Rampilov M.O., Ripp G.S., Lastochkin E.I., Khubanov V.B., Posokhov V.F., Vladyskin N.V. Age and mineralogical and geochemical parameters of rocks of the China alkaline massif (western Transbaikalia) // *Russian Geology and Geophysics*. 2017, 58, pp. 903–921.
- Izokh A.E., Polyakov G.V., Shelepaev R.A., Vrublevskii V.V., Egorova V.V., Rudnev S.N., Lavrenchuk A.V., Borodina E.V., Oyunchimeg T. Early Paleozoic Large Igneous Province of the Central Asia Mobile Belt. <http://www.largeigneousprovinces.org/08may>. May 2008 LIP of the Month.
- Jahn B.-M., Wu F.Y., Chen B. Massive granitoid generation in Central Asia: Nd isotope evidence and implication for continental growth in the Phanerozoic // *Episodes*. 2000, 23, pp. 82–92.
- Kelemen P.B., Hanghøj K., Greene A.R. One view of the geochemistry of subduction-related magmatic arcs, with an emphasis on primitive andesite and lower crust. In *Treatise on Geochemistry*, Eds. Holland Y.D., Turekian K.K., Elsevier Ltd.: Amsterdam, 2003, 3, pp. 593–659.
- Khain E.V., Bibikova E.V., Salnikova E.B., Kröner A., Gibsher A.S., Didenko A.N., Degtyarev K.E., Fedotova A.A. The Palaeo-Asian Ocean in the Neoproterozoic and Early Palaeozoic: new geochronologic data and palaeotectonic reconstructions // *Precambrian Research*. 2003, 122, pp. 329–358.
- Kheraskova T.N., Bush V.A., Didenko A.N., Samygin S.G. Breakup of Rodinia and early stages of evolution of the Paleozoic Ocean // *Geotectonics*. 2010, 44, pp. 3–24.
- Krupchatnikov V.I., Vrublevskii V.V., Kruk N.N. Early Devonian volcanics of southeastern Gorny Altai: geochemistry, isotope (Sr, Nd, and O) composition, and petrogenesis (Aksai complex) // *Russian Geology and Geophysics*. 2018, 59, pp. 905–924.
- Kungurtsev L.V., Berzin N.A., Kazanskii A.Yu., Metelkin D.V. Tectonic evolution of the southwestern framing of the Siberian Platform in the Vendian–Cambrian according to paleomagnetic data // *Russian Geology and Geophysics*. 2001, 42, pp. 1042–1051.
- Kuzmin M.I., Yarmolyuk V.V., Kravchinsky V.A. Phanerozoic hot spot traces and paleogeographic reconstructions of the Siberian continent based on interaction with the African large low shear velocity province // *Earth-Science Reviews*. 2010, 102, pp. 29–59.
- Li Z.X., Bogdanova S.V., Collins A.S., Davidson A., De Waele B., Ernst R.E., Fitzsimons I.C.W., Fuck R.A., Gladkochub D.P., Jacobs J., Karlstrom K.E., Lu S., Natapov L.M., Pease V., Pisarevsky S.A., Thrane K., Vernikovskiy V. Assembly, configuration, and break-up history of Rodinia: A synthesis // *Precambrian Research*. 2008, 160, pp. 179–210.
- Lightfoot P.C., Hawkesworth C.J., Hergt J., Naldrett A.J., Gorbachev N.S., Fedorenko V.A., Doherty W. Remobilisation of the major, trace-element, and from picritic and tholeiitic Siberian Trap, Russia // *Contributions to Mineralogy and Petrology*. 1993, 114, pp. 171–188, DOI 10.1007/BF00307754.
- Mustafaev A.A., Gertner I.F., Ernst R.E., Serov P.A., Kolmakov Y.V. The Paleozoic-aged University foidolite-gabbro pluton of the northeastern part of the Kuznetsk Alatau Ridge, Siberia: Geochemical characterization, geochronology, petrography and geophysical indication of potential high-grade nepheline ore // *Minerals*. 2020, 10, 1128. <https://doi.org/10.3390/min10121128>.
- Nikiforov A.V., Yarmolyuk V.V. Early Paleozoic age and geodynamic setting of the Botogol and Khushagol alkaline massifs in the Central Asian fold belt // *Doklady Earth Sciences*. 2007, 412, pp. 6–10, doi:10.1134/S1028334X07010023.
- Nikiforov A.V., Yarmolyuk V.V., Kovalenko V.I., Ivanov V.G., Zhuravlev D.Z. Late Mesozoic carbonatites of western Transbaikalia: Isotopic–geochemical characteristics and sources // *Petrology*. 2002, 10, pp. 146–164.
- Pearce J.A., Harris N.B.W., Tindle A.G. Trace element discrimination diagrams for the tectonic interpretation of granitic rocks // *Journal of Petrology*. 1984, 25, pp. 956–983, doi:10.1093/petrology/25.4.956.
- Pokrovskii B.G., Andreeva E.D., Vrublevskii V.V., Grinev O.M. Contamination mechanisms of alkaline-gabbro intrusions in the southern periphery of the Siberian craton: evidence from strontium and oxygen isotopic compositions // *Petrology*. 1998, 6, pp. 237–251.
- Rudnev S.N. *Rannepaleozoyskiy granitoidnyy magmatizm Altaye-Sayanskoy skladchatoy oblasti i Ozerney zony Zapadnoy Mongolii* [Early Paleozoic Granitoid Magmatism in the Altai-Sayan Folded Area and in the Lake Zone in Western Mongolia] Novosibirsk: Publ. House SB RAS, 2013, 300 p. In Russian: Руднев С.Н. Раннепалеозойский гранитоидный магматизм Алтае-Саянской складчатой области и Озерной зоны Западной Монголии. Новосибирск, Изд-во СО РАН, 2013, 300 с.

- Rudnick R.L., Gao S. Composition of the continental crust, Treatise on Geochemistry In Treatise on Geochemistry; Eds. Holland Y.D., Turekian K.K.; Amsterdam: Elsevier Ltd., 2003, 3, pp. 1–50.
- Saccani E. A new method of discriminating different types of post-Archean ophiolitic basalts and their tectonic significance using Th–Nb and Ce–Dy–Yb systematics // *Geoscience Frontiers*. 2015, 6, pp. 481–501.
- Salnikova E.B., Stifeeva M.V., Nikiforov A.V., Yarmolyuk V.V., Kotov A.B., Anisimova I.V., Sugorakova A.M., Vrublevskii V.V. Andradite–morimotoite garnets as promising U–Pb geochronometers for dating ultrabasic alkaline rocks // *Doklady Earth Sciences*. 2018, 480 (2), pp. 778–782.
- Salters V., Stracke A. Composition of the depleted mantle. *Geochemistry, Geophysics, Geosystems* 2004, 5, C00597.
- Şengör A.C., Natal'in B.A., Burtman V.S. Evolution of the Altaid tectonic collage and Palaeozoic crustal growth in Eurasia // *Nature*. 1993, 364, pp. 299–306.
- Sklyarov E.V., Fedorovsky V.S., Kotov A.B., Lavrenchuk A.V., Mazukabzov A.M., Levitsky V.I., Sal'nikova E.B., Starikova A.E., Yakovleva S.Z., Anisimova I.V., Fedoseenko A.M. Carbonatites in collisional settings and pseudo-carbonatites of the Early Paleozoic Ol'khon collisional system // *Russian Geology and Geophysics* 2009, 50, pp. 1091–1106.
- Stracke A., Hofmann A.W., Hart S.R. FOZO, HIMU, and the rest of the mantle zoo. *Geochemistry, Geophysics, Geosystems* 2005, 6, Q05007, DOI 10.1029/2004GC000824
- Sun S., McDonough W.F. Chemical and isotopic systematics of oceanic basalts: implications for mantle composition and processes. In *Magmatism in the Ocean Basins*, Eds. Saunders A.D., Norry M.J., Geol. Soc. Spec. Publ., 1989, 42, pp. 313–345.
- Thiéblemont D. Discrimination between mantle- and crustal-derived Andean calc-alkaline lavas. *Paris: C.R. Acad. Sci.* 1999, 329, pp. 243–250.
- Vorontsov A., Yarmolyuk V., Dril S., Ernst R., Perfilova O., Grinev O., Komaritsyna T. Magmatism of the Devonian Altai-Sayan Rift System: Geological and geochemical evidence for diverse plume-lithosphere interactions // *Gondwana Research*. 2021, 89, pp. 193–219.
- Vrublevskii V.V., Gertner I.F., Zhuravlev D.Z., Makarenko N.A. Sm–Nd isotopic age and nature of source for the association of alkaline basic rocks and carbonatites of the Kuznetskii Alatau // *Doklady Akademii Nauk*. 2003, 391(3), pp. 378–383.
- Vrublevskii V.V., Voitenko N.N., Romanov A.P., Polyakov G.V., Izokh A.E., Gertner I.F., Krupchatnikov V.I. Magma sources of Triassic lamproites of Gornyi Altai and Taimyr: Sr and Nd isotope evidence for plume-lithosphere interaction // *Doklady Earth Sciences*. 2005, 405A, pp. 1365–1367.
- Vrublevskii V.V., Krupchatnikov V.I., Izokh A.E., Gertner, I.F. The alkaline and carbonatitic rocks of Gornyi Altai (Edel'veis complex) as indicators of Early Paleozoic plume magmatism in the Central Asian Fold Belt // *Russian Geology and Geophysics*. 2012, 53, pp. 721–735.
- Vrublevskii V.V., Gertner I.F., Gutiérrez-Alonso G., Hofmann M., Grinev O.M., Tishin P.A. Isotope (U–Pb, Sm–Nd, Rb–Sr) geochronology of alkaline basic plutons of the Kuznetsk Alatau // *Russian Geology and Geophysics*. 2014, 55, pp. 1264–1277.
- Vrublevskii V.V. Sources and geodynamic setting of petrogenesis of the Middle Cambrian Upper Petropavlovka alkaline basic pluton (Kuznetsk Alatau, Siberia) // *Russian Geology and Geophysics*. 2015, 56, 379–401.
- Vrublevskii V.V., Kotelnikov A.D., Rudnev S.N., Krupchatnikov V.I. Evolution of the Paleozoic granitoid magmatism in the Kuznetsk Alatau: new geochemical and U–Pb (SHRIMP-II) isotope data // *Russian Geology and Geophysics*. 2016a, 57, pp. 225–246.
- Vrublevskii V.V., Grinev O.M., Izokh A.E., Travin A.V. Geochemistry, isotope triad (Nd–Sr–O), and ^{40}Ar – ^{39}Ar age of Paleozoic alkaline mafic intrusions of the Kuznetsk Alatau (by the example of the Belaya Gora pluton) // *Russian Geology and Geophysics*. 2016b, 57, pp. 592–602.
- Vrublevskii V.V., Gertner I.F., Chugaev A.V. Parental sources of high-alumina alkaline melts: Nd, Sr, Pb, and O isotopic evidence from the Devonian Kiya–Shaltyr gabbro–urtite intrusion, South Siberia // *Doklady Earth Sciences*. 2018, 479 (2), pp. 518–523.
- Vrublevskii V.V., Kotelnikov A.D., Izokh A.E. The age and petrologic and geochemical conditions of formation of the Kogtakht gabbro–monzonite complex in the Kuznetsk Alatau // *Russian Geology and Geophysics*. 2018a, 59, pp. 718–744.
- Vrublevskii V.V., Morova A.A., Bukharova O.V., Konovalenko, S.I. Mineralogy and geochemistry of Triassic carbonatites in the Matcha alkaline intrusive complex (Turkestan–Alai Ridge, Kyrgyz Southern Tien Shan), SW Central Asian orogenic belt // *Journal of Asian Earth Sciences*. 2018b, 153, pp. 252–281.
- Vrublevskii V.V., Gertner I.F., Ernst R.E., Izokh A.E., Vishnevskii A.V. The Overmarat–Gol alkaline pluton in Northern Mongolia: U–Pb age and preliminary implications for magma sources and tectonic setting // *Minerals*. 2019, 9, 170, doi.org/10.3390/min9030170.
- Vrublevskii V.V., Nikiforov A.V., Sugorakova A.M., Kozulina T.V. Petrogenesis and tectonic setting of the Cambrian Kharly alkaline–carbonatite complex (Sangilen Plateau, Southern Siberia): implications for the Early Paleozoic evolution of magmatism in the western Central Asian Orogenic Belt // *Journal of Asian Earth Sciences*. 2020a, 188, 104163, DOI: 10.1016/j.jseas.2019.104163.
- Vrublevskii V.V., Gertner I.F., Gutiérrez-Alonso G., Hofmann M., Grinev O.M., Mustafayev A. Multiple intrusion stages and mantle sources of the Paleozoic Kuznetsk Alatau alkaline province, Southern Siberia: geochemistry and Permian U–Pb, Sm–Nd ages in the Goryachegorsk ijolite–foyaite intrusion // *International Geology Review*. 2020b, doi.org/10.1080/00206814.2020.1830312.
- Vrublevskii V.V., Gertner I.F. Paleozoic alkaline–mafic intrusions of the Kuznetsk Alatau, their sources and conditions for magma generation // *Petrology*. 2021, 29, pp. 24–53.
- Wan B., Li Sh., Xiao W., Windley B.F. Where and when did the Paleo-Asian Ocean form? // *Precambrian Research*. 2018, 317, pp. 241–252.
- Wilhelm C., Windley B.F. Tectonic synthesis of the altaids of Central Asia. In *The Central Asian Orogenic Belt: Geology, Evolution, Tectonics and Models*; Ed. Kröner, A.; Borntraeger Science Publishers: Stuttgart, 2015, pp. 235–252.
- Windley B.F., Alexeiev D., Xiao W.J., Kröner A., Badarch G., Tectonic models for accretion of the Central Asian orogenic belt // *J. Geol. Soc. London* 2007, 164, pp. 31–47.
- Yarmolyuk V.V., Kovalenko V.I. Deep geodynamics and mantle plumes: their role in the formation of the Central Asian orogenic belt. *Petrology*. 2003, 11, pp. 504–531.
- Yarmolyuk V.V., Kovalenko V.I., Kovach, V.P. Kozakov, I.K., Kotov, A.B.; Sal'nikova, E.B. Geodynamics of Caledonides in the Central Asian fold belt // *Doklady Earth Sciences*. 2003, 389A, pp. 311–316.

Yarmolyuk V.V., Kuzmin M.I., Ernst R.E. Intraplate geodynamics and magmatism in the evolution of the Central Asian Orogenic Belt // *Journal of Asian Earth Sciences*. 2014, 93, pp. 158–179.

Yarmolyuk V.V., Nikiforov A.V., Sal'nikova E.B., Travin A.V., Kozlovskiy A.M., Kotov A.B., Shuriga T.N., Lyukhin D.A., Lebedev V.I., Anisimova I.V., Plotkina Yu.V., Yakovleva S.Z. Rare-metal granitoids of the Ulug-Tanzek deposit (Eastern Tyva): age and tectonic setting // *Doklady Earth Sciences*. 2010, 430(1), pp. 95–100.

Yashina R.M. *Shchelochnoy magmatizm skladchato-glybovykh oblastey* [Alkaline Magmatism in Orogenic Areas] Moscow: Nauka, 1982, 274 p. In Russian: Яшина Р.М. *Щелочной магматизм складчато-глубоких областей*. М.: Наука, 1982. 274 с.

Zindler A., Hart S.R. Chemical geodynamics. *Annual Review of Earth and Planetary Sciences*. 1986. 14. pp. 493–571.

Information about the author:

Vrublevskii V.V., Dr. Sci. (Geol.-Miner.), Head of Department of Dynamical Geology, Geology and Geography Faculty, National Research Tomsk State University, Tomsk, Russia.

E-mail: vasvr@yandex.ru

The author declares no conflicts of interests.

Информация об авторах:

Врублевский В.В., доктор геолого-минералогических наук, заведующий кафедрой динамической геологии, геолого-географический факультет, Национальный исследовательский Томский государственный университет, Томск, Россия.

E-mail: vasvr@yandex.ru

Автор заявляет об отсутствии конфликта интересов.

The article was submitted 25.09.2021; approved after reviewing 28.11.2021; accepted for publication 12.09.2022

Статья поступила в редакцию 25.09.2021; одобрена после рецензирования 28.11.2021; принята к публикации 12.09.2022